



## Invasive Exotic Plant Monitoring at Lincoln Boyhood National Memorial: Year 1 (2006)

Natural Resource Technical Report NPS/HTLN/NRTR—2007/021  
NPS D-43



**ON THE COVER**

Invasive, exotic plants at Lincoln Boyhood National Memorial. Left to right: A) Japanese honeysuckle. B) Periwinkle near visitor's center. C) Periwinkle in forest. D) Chinese yam shading ground cover. E) Chinese yam growing into the canopy. F) Fruiting Chinese yam. G) Typical privet shrub. H) Park staff standing amidst periwinkle and privet. I) Multiflora rose seedbank. Photographs by Michael H. Williams.

---

**Invasive Exotic Plant Monitoring at Lincoln Boyhood National Memorial:  
Year 1 (2006)**

Natural Resource Technical Report NPS/HTLN/NRTR—2007/021  
NPS D-43

J. Tyler Cribbs, Craig C. Young, Jennifer L. Haack, and Holly J. Etheridge  
National Park Service, Heartland I&M Network  
Wilson's Creek National Battlefield, 6424 West Farm Road 182, Republic, MO 65738



March 2007

U.S. Department of the Interior  
National Park Service  
Natural Resource Program Center  
Fort Collins, Colorado

The Natural Resource Publication series addresses natural resource topics that are of interest and applicability to a broad readership in the National Park Service and to others in the management of natural resources, including the scientific community, the public, and the NPS conservation and environmental constituencies. Manuscripts are peer-reviewed to ensure that the information is scientifically credible, technically accurate, appropriately written for the intended audience, and is designed and published in a professional manner.

The Natural Resource Technical Report series is used to disseminate the peer-reviewed results of scientific studies in the physical, biological, and social sciences for both the advancement of science and the achievement of the National Park Service's mission. The reports provide contributors with a forum for displaying comprehensive data that are often deleted from journals because of page limitations. Current examples of such reports include the results of research that addresses natural resource management issues; natural resource inventory and monitoring activities; resource assessment reports; scientific literature reviews; and peer reviewed proceedings of technical workshops, conferences, or symposia.

Views and conclusions in this report are those of the authors and do not necessarily reflect policies of the National Park Service. Mention of trade names or commercial products does not constitute endorsement or recommendation for use by the National Park Service.

Printed copies of reports in these series may be produced in a limited quantity and they are only available as long as the supply lasts. This report is also available from the Heartland I&M Network website (<http://www.nature.nps.gov/im/units/HTLN>) on the internet, or by sending a request to the address on the back cover.

Please cite this publication as:

Cribbs, J.T., C.C. Young, J.L. Haack, and H.J. Etheridge. 2007. Invasive exotic plant monitoring at Lincoln Boyhood National Memorial: Year 1 (2006). Natural Resource Technical Report NPS/HTLN/NRTR—2007/021. National Park Service, Fort Collins, Colorado.

NPS D-43, March 2007

## Acknowledgements

Dan Tenaglia conducted all field work associated with this report in 2006. Dan died on February 13, 2007 following a collision two days earlier while riding his bike near his home in Opelika, Alabama. We are grateful for his work. Dan's botanical photography can be found at his Missouri Plants website: <http://www.missouriplants.com/>.

## **Executive Summary**

During surveys in 2006, we documented 31 invasive exotic plant taxa at Lincoln Boyhood National Memorial. Japanese honeysuckle was widespread and abundant at Lincoln Boyhood National Memorial. Japanese honeysuckle was established on at least 3.2 acres and occurred in 94.8% of search units. Common periwinkle was the second most abundant species with a cover of 1.3 acres and occurred in 24.7% of search units. A total of 27 invasive exotic plant species each covered one acre or less of the memorial. In general, several invasive exotic plants are a moderate problem at Lincoln Boyhood National Memorial, but successful control appears possible for a large group of species. The acreage estimates presented in the report may be used to plan management activities leading to control of exotic plants and the accomplishment of GPRA goal IA1b.

# Table of Contents

	Page
Introduction.....	1
Methods.....	3
Results and Discussion .....	4
Literature Cited.....	4

This page intentionally left blank.

## Introduction

*Author's note.* In this report, we use the term invasive exotic plant to refer to plants that are not native to the park and that are presumed to pose environmental harm to native plant populations and/or communities based on a review of numerous state and regional invasive exotic plant lists. The great majority of the introductory text was taken from Welch and Geissler (2007) with slight modification.

**Scope of invasive exotic plant problem for National Parks.** Globalization of commerce, transportation, human migration, and recreation in recent history has introduced invasive exotic species to new areas at an unprecedented rate. Biogeographical barriers that once restricted the location and expansion of species have been circumvented, culminating in the homogenization of the Earth's biota. Although only 10% of introduced species become established and only 1% become problematic (Williamson 1993, Williamson and Fitter 1996) or invasive, nonnative species have profound impacts worldwide on the environment, economies, and human health. Invasive species have been directly linked to the replacement of dominant native species (Tilman 1999), the loss of rare species (King 1985), changes in ecosystem structure, alteration of nutrient cycles and soil chemistry (Ehrenfeld 2003), shifts in community productivity (Vitousek 1990), reduced agricultural productivity, and changes in water availability (D'Antonio and Mahall 1991). Often the damage caused by these species to natural resources is irreparable and our understanding of the consequences incomplete. Invasive species are second only to habitat destruction as a threat to wildland biodiversity (Wilcove et al. 1998). Consequently, the dynamic relationships among plants, animals, soil, and water established over many thousands of years are at risk of being destroyed in a relatively brief period.

For the National Park Service (NPS), the consequences of these invasions present a significant challenge to the management of the agency's natural resources "unimpaired for the enjoyment of future generations." National Parks, like other land management organizations, are deluged by new exotic species arriving through predictable (e.g., road, trail, and riparian corridors), sudden (e.g., long-distance dispersal through cargo containers and air freight), and unexpected anthropogenic pathways (e.g., weed seeds in restoration planting mixes). Nonnative plants claim an estimated 4,600 acres of public lands each year in the United States (Asher and Harmon 1995), significantly altering local flora. For example, exotic plants comprise an estimated 43% and 36% of the flora of the states of Hawaii and New York, respectively (Rejmanek and Randall 1994). Invasive plants infest an estimated 2.6 million acres of the 83 million acres managed by the NPS.

More NPS lands are infested daily despite diligent efforts to curtail the problem. Impacts from invasive species have been realized in most parks, resulting in an expressed need to control existing infestations and restore affected ecosystems. Additionally, there is a growing urgency to be proactive—to protect resources not yet impacted by current and future invasive species (Marler 1998). Invasive exotic species most certainly will continue to be a management priority for the National Parks well into the 21st Century. Invasive exotic plants have been consistently ranked as a top vital sign for long term monitoring as part of the NPS Inventory & Monitoring (I&M) Program. During the vital signs selection process in 2003, Heartland Network parks recognized the need for exotic plant monitoring (DeBacker et al. 2004). Nine parks (CUVA,

EFMO, GWCA, HEHO, HOCU, HOME, LIBO, OZAR, PERI) identified invasive exotic plants as their most important management issue, two parks (TAPR, WICR) identified invasive exotic plants as their second most important management issue, and PIPE identified invasive exotic plants as its third most important management issue. During this process, invasive exotic plant monitoring was recognized across all network parks as the most important shared monitoring need.

**Prevention and early detection as keys to invasive exotic plant management.** Prevention and early detection are the principal strategies for successful invasive exotic plant management. While there is a need for long-term suppression programs to address very high-impact species, eradication efforts are most successful for infestations less than one hectare in size (Rejmanek and Pitcairn 2002). Eradication of infestations larger than 100 hectares is largely unsuccessful, costly, and unsustainable (Rejmanek and Pitcairn 2002). Costs, or impacts, to ecosystem components and processes resulting from invasion also increase dramatically over time, making ecosystem restoration improbable in the later stages of invasion. Further, in their detailed review of the nonnative species problem in the United States, the US Congress, Office of Technology Assessment (1993) stated that the environmental and economic benefits of supporting prevention and early detection initiatives significantly outweigh any incurred costs, with the median benefit-to-cost ratio being 17:1 in favor of being proactive.

Although preventing the introduction of invasive exotic plants is the most successful and preferred strategy for resource managers, the realities of globalization, tight fiscal constraints, and limited staff time guarantee that invaders will get through park borders. Fortunately, invasive exotic plants quite often undergo a lag period between introduction and subsequent colonization of new areas. Managers, then, can take advantage of early detection monitoring to make certain invasive exotic species are found and successfully eradicated before populations become well established.

This strategy requires resource managers to: (1) detect invasive exotic species early (i.e., find a new species or an incipient population of an existing species while the infestation is small (less than 1 hectare), and (2) respond rapidly (i.e., implement appropriate management techniques to eliminate the invasive plant and all of its associated regenerative material).

**Invasive exotic plant management at Lincoln Boyhood National Memorial.** While a complete history of park invasive exotic plant management issues is beyond the scope of this report, a few important highlights are given:

1. The forest at Lincoln Boyhood National Memorial is a significant cultural resource that is vulnerable to exotic plant invasions.
2. Based on a 2003 survey, the following invasive, exotic plants were found at Lincoln Boyhood National Memorial: multiflora rose (*Rosa multiflora*), privet species (*Ligustrum* spp.), common periwinkle (*Vinca minor*), sericea lespedeza (*Lespedeza cuneata*), and Chinese yam (*Dioscorea oppositifolia*).

3. Park use of mechanical and chemical methods of control will be needed to control a number of invasive, exotic plant species.

## Methods

**Watch lists.** The invasive exotic plants on three watch lists were sought during monitoring (Table 1). Invasive exotic plants not known to occur on the park based on NPSpecies (the national NPS database for plant occurrence registration) constitute the early detection watch list. Invasive exotic plants known to occur on the park based on NPSpecies constitute the park-established watch list. Invasive exotic plants from the park-based watch list included plants selected by park managers or network staff which may not have been included on the other lists due to incomplete information in NPSpecies (e.g., not documented) or USDA Plants (e.g., state distribution information inaccurate) databases or due to differing opinions regarding network designation of a plant as a high priority. While aquatic species are listed on the watch lists, terrestrial plants were the focus of this survey.

**Field methods.** Invasive exotic plant species on designated watch lists (Table 1) were sought in high priority areas on Lincoln Boyhood National Memorial (Figure 1). Dan Tenaglia, the contract botanist for this project navigated through search units using a Thales GPS unit, identified invasive exotic plants in an approximately 6-m belt, and attributed a coarse cover value to each species (0=0, 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9 m<sup>2</sup>). A total of 77 search units were surveyed at Lincoln Boyhood National Memorial. The botanist had discretion to search a larger belt if feasible, to target locations likely to support exotic plants (e.g., field edges, roads), and to circumvent extremely difficult or hazardous terrain when needed. Cover was estimated for all plants observed while navigating in the search unit (i.e., not restricted to the 6-m belt).

**Analytical methods.** Data analysis involved simple displays, as well as calculation of plant frequency and cover. The invasive exotic plants encountered on Lincoln Boyhood National Memorial were attributed to search units in a GIS (Figures 2 – 32). Note that entire search units were not fully searched. A park-wide cover range was estimated using the high and low values of the cover classes for each invasive exotic plant encountered, assuming that 20 % of the park was searched and that the areas searched were representative of the entire park. The park-wide frequency of invasive exotic plants was calculated as the percentage of occupied search units.

**Invasiveness ranks.** In order to provide additional information on the ecological impact and feasibility of control, the ecological impact and general management difficulty sub-ranks that constitute the invasiveness rank (I-rank), as determined by NatureServe (Morse et al. 2004), were listed when available. The ecological impact characterizes the effect of the plant on ecosystem processes, community composition and structure, native plant and animal populations, and the conservation significance of threatened biodiversity. General management difficulty ranks are assigned based on the resources and time generally required to control a plant, the non-target effects of control on native populations, and the accessibility of invaded sites. Sub-ranks are given as high (H), medium (M), low (L), insignificant (I), unknown (U), or a combination of ranks.

## Results and Discussion

In 2006, a total of 31 invasive exotic plant taxa were found during the survey at Lincoln Boyhood National Memorial (Table 2). Of these, 17 species were listed in NPSpecies as occurring at the memorial. Eleven species and three species were identified from the early detection watch list and park based watch lists, respectively.

The distribution and abundance of the invasive exotic plant species at Lincoln Boyhood National Memorial varied widely. Japanese honeysuckle (*Lonicera japonica*) was the most abundant and widespread invasive exotic plant species on the park. Japanese honeysuckle was found in 94.8 % of search units and was estimated to cover between three and ten acres. The next three most abundant species included common periwinkle (*Vinca minor*), European privet (*Ligustrum vulgare*), and multiflora rose (*Rosa multiflora*), covering at least 1.3, 0.7, and 0.6 acres, respectively. The remaining 27 invasive exotic plant species each covered less than one acre of the park.

Only two species were noted as having definitively high ecological impacts: autumn olive (*Elaeagnus umbellata*) and crownvetch (*Securigera varia*) (Table 2). Nine species were characterized as having at least a medium ecological impact. The remaining species had medium-low ecological impacts or less, including six species with low or insignificant impacts. Recognizing that the feasibility of control often strongly influences decisions regarding invasive exotic plant management, autumn olive, crownvetch, and Japanese barberry (*Berberis thunbergii*) with potentially high ecological impacts were noted as having low or insignificant management difficulty. Controlling these species will likely provide a high benefit for the management costs. Furthermore, management of the numerous small invasive exotic populations on Lincoln Boyhood National Memorial may well limit the financial and ecological costs associated with the spread of these species.

In summary, this report provides information on invasive, exotic plant abundance and distribution. The report also characterizes the ecological impacts and management difficulty associated with these species. The information is designed to assist park natural resource managers in planning invasive exotic plant management. The following links may further assist managers: <http://www.nature.nps.gov/im/units/htln/monitoring/projects/inp.htm> and <http://www.natureserve.org/explorer/>.

## Literature Cited

Asher, J. A., and D. W. Harmon. 1995. Invasive exotic plants are destroying the naturalness of U.S. Wilderness areas. *International Journal of Wilderness* 1:35-37.

D'Antonio, C. M., and B. E. Mahall. 1991. Root profiles and competition between the invasive, exotic perennial, *Carpobrotus edulis*, and two native shrub species in California coastal scrub. *American Journal of Botany* 78:885-894.

DeBacker, M.D., C.C. Young (editor), P. Adams, L. Morrison, D. Peitz, G.A. Rowell, M. Williams, and D. Bowles. 2005. Heartland Inventory and Monitoring Network and Prairie

Cluster Prototype Monitoring Program Vital Signs Monitoring Plan. National Park Service, Heartland Inventory and Monitoring Network and Prairie Cluster Prototype Monitoring Program, Wilson's Creek National Battlefield, Republic, Missouri, 104 pp. plus appendices.

Ehrenfeld, J.G. 2003. The effects of exotic plant invasions on soil nutrient cycling processes. *Ecosystems* 6:503-523.

King, W. B. 1985. Island birds: will the future repeat the past? Pages 3-15 in P. J. Moors, editor. *Conservation of Island Birds*. International Council for Bird Preservation. Cambridge University Press, Cambridge, UK.

Marler, M. 1998. Exotic plant invasions of federal Wilderness areas: current status and future directions. The Aldo Leopold Wilderness Research Institute. Rocky Mountain Research Station, Missoula, Montana, USA.

Office of Technology Assessment. 1993. Harmful non-indigenous species in the United States. OTA-F-565. U.S. Congress, Government Printing Office, Washington, D.C., USA.

Rejmanek, M., and M. J. Pitcairn. 2002. When is eradication of exotic pest plants a realistic goal? Pages 249-253 in C. R. Veitch and M. N. Clout, editors. *Turning the Tide: the Eradication of Invasive Species*. IUCN SSC Invasive Species Specialist Group. IUCN, Gland, Switzerland and Cambridge, UK.

Rejmanek, M., and J. M. Randall. 1994. Invasive alien plants in California: 1993 summary and comparison with other areas in North America. *Madrono* 41:161-177.

Tilman, D. 1999. The ecological consequences of changes in biodiversity: a search for general principles. *Ecology* 80:1455-1474.

Vitousek, P. M. 1990. Biological invasions and ecosystem processes: towards an integration of population biology and ecosystem studies. *Oikos* 57:7-13.


Welch, B.A. and P.H. Geissler. 2007. Early detection of invasive plants: a handbook. United States Geological Survey draft. <http://www.pwrc.usgs.gov/brd/invasiveHandbook.cfm>.

Wilcove, D. S., D. Rothstein, J. Dubow, A. Phillips, and E. Losos. 1998. Quantifying threats to imperiled species in the United States. *Bioscience* 48:607-615.

Williamson, M. 1993. Invaders, weeds and risk from genetically modified organisms. *Experientia* 49:219-224.

Williamson, M. and A. Fitter. 1996. The varying success of invaders. *Ecology* 77:1661-1666.

# Lincoln Boyhood National Memorial Exotic Plant Search Units



**Figure 1. Invasive exotic plant search units at Lincoln Boyhood National Memorial. The search units indicate the search locations for invasive exotic plants in 2006.**

**Table 1. Watch lists for Lincoln Boyhood National Memorial**

Early Detection Watch List		Park-Established Watch List		Park-Based Watch List	
<i>Acer platanoides</i>	Norway maple	<i>Albizia julibrissin</i>	Silktree	<i>Ligustrum obtusifolium</i>	Border privet
<i>Ailanthus altissima</i>	Tree of heaven	<i>Berberis thunbergii</i>	Japanese barberry	<i>Wisteria sinensis</i>	Chinese wisteria
<i>Alliaria petiolata</i>	Garlic mustard	<i>Dactylis glomerata</i>	Orchardgrass	<i>Ligustrum sinense</i>	Chinese privet
<i>Alnus glutinosa</i>	European alder	<i>Dioscorea oppositifolia</i>	Chinese yam		
<i>Arctium minus</i>	Lesser burdock	<i>Euonymus alatus</i>	Burningbush		
<i>Azolla</i>	Mosquitofern	<i>Euonymus fortunei</i>	Winter creeper		
<i>Bromus inermis</i>	Smooth brome	<i>Glechoma hederacea</i>	Ground ivy		
<i>Bromus sterilis</i>	Poverty brome	<i>Hedera helix</i>	English ivy		
<i>Bromus tectorum</i>	Cheatgrass	<i>Lespedeza cuneata</i>	Sericea lespedeza		
<i>Butomus umbellatus</i>	Flowering rush	<i>Ligustrum vulgare</i>	European privet		
<i>Carduus nutans</i>	Nodding plumeless thistle	<i>Lolium pratense</i>	Meadow fescue		
<i>Celastrus orbiculatus</i>	Oriental bittersweet	<i>Lonicera japonica</i>	Japanese honeysuckle		
<i>Centaurea biebersteinii</i>	Spotted knapweed	<i>Lonicera morrowii</i>	Morrow's honeysuckle		
<i>Centaurea solstitialis</i>	Yellow star-thistle	<i>Melilotus officinalis</i>	Yellow sweetclover		
<i>Cirsium arvense</i>	Canada thistle	<i>Morus alba</i>	White mulberry		
<i>Cirsium vulgare</i>	Bull thistle	<i>Plantago lanceolata</i>	Narrowleaf plantain		
<i>Cynanchum louiseae</i>	Louise's swallow-wort	<i>Poa compressa</i>	Canada bluegrass		
<i>Cynanchum rossicum</i>	European swallow-wort	<i>Poa pratensis</i>	Kentucky bluegrass		
<i>Dipsacus fullonum</i>	Fuller's teasel	<i>Polygonum cuspidatum</i>	Japanese knotweed		
<i>Dipsacus laciniatus</i>	Cutleaf teasel	<i>Robinia pseudoacacia</i>	Black locust		
<i>Egeria densa</i>	Brazilian waterweed	<i>Rosa multiflora</i>	Multiflora rose		
<i>Elaeagnus umbellata</i>	Autumn olive	<i>Verbascum thapsus</i>	Common mullein		
<i>Euphorbia esula</i>	Leafy spurge	<i>Vinca minor</i>	Common periwinkle		
<i>Frangula alnus</i>	Glossy buckthorn				
<i>Hesperis matronalis</i>	Dames rocket				
<i>Holcus lanatus</i>	Common velvetgrass				
<i>Humulus japonicus</i>	Japanese hop				
<i>Hyoscyamus niger</i>	Black henbane				
<i>Lespedeza bicolor</i>	Shrub lespedeza				
<i>Lolium arundinaceum</i>	Tall fescue				
<i>Lonicera maackii</i>	Amur honeysuckle				
<i>Lonicera tatarica</i>	Tatarian honeysuckle				
<i>Lonicera X bella</i>	Showy fly honeysuckle				
<i>Lotus corniculatus</i>	Bird's-foot trefoil				
<i>Lysimachia nummularia</i>	Creeping jenny				
<i>Lythrum salicaria</i>	Purple loosestrife				


**Table 1. Watch lists for Lincoln Boyhood National Memorial (cont.)**

Early Detection Watch List		Park-Established Watch List		Park-Based Watch List	
<i>Microstegium vimineum</i>	Nepalese browntop				
<i>Miscanthus sinensis</i>	Chinese silvergrass				
<i>Myriophyllum spicatum</i>	Eurasian watermilfoil				
<i>Ornithogalum umbellatum</i>	Sleepydick				
<i>Pastinaca sativa</i>	Wild parsnip				
<i>Paulownia tomentosa</i>	Princesstree				
<i>Phalaris arundinacea</i>	Reed canarygrass				
<i>Phragmites australis</i>	Common reed				
<i>Populus alba</i>	White poplar				
<i>Potamogeton crispus</i>	Curly pondweed				
<i>Potentilla recta</i>	Sulphur cinquefoil				
<i>Pueraria montana var. lobata</i>	Kudzu				
<i>Rhamnus cathartica</i>	Common buckthorn				
<i>Securigera varia</i>	Crownvetch				
<i>Solanum dulcamara</i>	Climbing nightshade				
<i>Sorghum halepense</i>	Johnsongrass				
<i>Spiraea japonica</i>	Japanese meadowsweet				
<i>Torilis arvensis</i>	Spreading hedgeparsley				
<i>Torillis japonica</i>	Erect hedgeparsley				
<i>Typha angustifolia</i>	Narrowleaf cattail				
<i>Ulmus pumila</i>	Siberian elm				
<i>Viburnum opulus</i>	European cranberrybush				

**Table 2. Overview of invasive exotic plants found on Lincoln Boyhood National Memorial. Ecological impact and general management difficulty based on NatureServe I-Rank subranks, Morse et al. 2004. Subranks are given as high (H), medium (M), low (L), insignificant (I), unknown (U), a range of ranks (indicated by /), or not available (--).**


Species	Common Name	Watch list	Park-wide cover (acres)	Frequency (percent)	Ecological impact	Management difficulty
<i>Lonicera japonica</i>	Japanese honeysuckle	Park-established	3.2 - 9.8	94.8%	M	HM
<i>Vinca minor</i>	Common periwinkle	Park-established	1.3 - 5.4	24.7%	I	U
<i>Ligustrum vulgare</i>	European privet	Park-established	0.7 - 2.7	68.8%	HL	HM
<i>Rosa multiflora</i>	Multiflora rose	Park-established	0.6 - 2.5	72.7%	L	L
<i>Robinia pseudoacacia</i>	Black locust	Park-established	< 1.0	32.5%	HM	M
<i>Ailanthus altissima</i>	Tree of heaven	Early detection	< 0.5	16.9%	ML	ML
<i>Elaeagnus umbellata</i>	Autumn olive	Early detection	< 0.5	28.6%	H	L
<i>Euonymus fortunei</i>	Winter creeper	Park-established	< 0.5	32.5%	M	LI
<i>Microstegium vimineum</i>	Nepalese browntop	Early detection	< 0.5	26.0%	M	HM
<i>Spiraea</i> spp	Meadowsweet	Early detection	< 0.5	5.2%	ML	HM
<i>Albizia julibrissin</i>	Silktree	Park-established	< 0.25	6.5%	ML	ML
<i>Celastrus orbiculatus</i>	Oriental bittersweet	Early detection	< 0.25	36.4%	ML	M
<i>Dioscorea oppositifolia</i>	Chinese yam	Park-established	< 0.25	9.1%	ML	MI
<i>Euonymus alatus</i>	Burningbush	Park-established	< 0.25	10.4%	LI	L
<i>Hedera helix</i>	English ivy	Park-established	< 0.25	6.5%	M	ML
<i>Lespedeza cuneata</i>	Sericea lespedeza	Park-established	< 0.25	14.3%	ML	ML
<i>Sorghum halepense</i>	Johnsongrass	Early detection	< 0.25	1.3%	ML	HM
<i>Dactylis glomerata</i>	Orchardgrass	Park-established	< 0.1	2.6%	LI	ML
<i>Ligustrum obtusifolium</i>	Border privet	Park-based	< 0.1	1.3%	LI	L
<i>Lonicera maackii</i>	Amur honeysuckle	Early detection	< 0.1	3.9%	HM	M
<i>Lonicera morrowii</i>	Morrow's honeysuckle	Park-established	< 0.1	3.9%	ML	M
<i>Lotus corniculatus</i>	Bird's-foot trefoil	Early detection	< 0.1	1.3%	ML	ML
<i>Lysimachia nummularia</i>	Creeping jenny	Early detection	< 0.1	9.1%	L	L
<i>Polygonum cuspidatum</i>	Japanese knotweed	Park-established	< 0.1	6.5%	HM	M
<i>Securigera varia</i>	Crownvetch	Early detection	< 0.1	2.6%	H	L
<i>Wisteria sinensis</i>	Chinese wisteria	Park-based	< 0.1	3.9%	ML	L
<i>Berberis thunbergii</i>	Japanese barberry	Park-established	< 0.01	9.1%	HM	I
<i>Cirsium vulgare</i>	Bull thistle	Early detection	< 0.01	9.1%	ML	ML
<i>Ligustrum sinense</i>	Chinese privet	Park-based	< 0.01	2.6%	M	L
<i>Poa compressa</i>	Canada bluegrass	Park-established	< 0.01	2.6%	ML	HL
<i>Verbascum thapsus</i>	Common mullein	Park-established	< 0.001	1.3%	ML	L

## *Ailanthus altissima*- 2006




**Figure 2. Abundance and distribution of *Ailanthus altissima* (tree of heaven) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1000-4999.9 m<sup>2</sup>, 8=5000-9999.9 m<sup>2</sup>, and 9=10,000-14999.9.**

### *Albizia julibrissin* - 2006




**Figure 3. Abundance and distribution of *Albizia julibrissin* (silktree) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Berberis thunbergii* - 2006




**Figure 4. Abundance and distribution of *Berberis thunbergii* (Japanese barberry) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Celastrus orbiculatus* - 2006




**Figure 5. Abundance and distribution of *Celastrus orbiculatus* (oriental bittersweet) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

### *Cirsium vulgare* - 2006




**Figure 6. Abundance and distribution of *Cirsium vulgare* (bull thistle) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Dactylis glomerata* - 2006




**Figure 7. Abundance and distribution of *Dactylis glomerata* (orchardgrass) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

*Dioscorea oppositifolia* - 2006




**Figure 8. Abundance and distribution of *Dioscorea oppositifolia* (chinese yam) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Elaeagnus umbellata* - 2006




**Figure 9. Abundance and distribution of *Elaeagnus umbellata* (autumn olive) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Euonymus alatus* - 2006




**Figure 10. Abundance and distribution of *Euonymus alatus* (burningbush) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Euonymus fortunei* - 2006




**Figure 11. Abundance and distribution of *Euonymus fortunei* (winter creeper) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Hedera helix* - 2006




**Figure 12. Abundance and distribution of *Hedera helix* (English ivy) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Lespedeza cuneata* - 2006




**Figure 13. Abundance and distribution of *Lespedeza cuneata* (*sericea lespedeza*) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

*Ligustrum obtusifolium* - 2006




**Figure 14. Abundance and distribution of *Ligustrum obtusifolium* (border privet) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Ligustrum sinense* - 2006




**Figure 15. Abundance and distribution of *Ligustrum sinense* (chinese privet) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Ligustrum vulgare* - 2006




**Figure 16. Abundance and distribution of *Ligustrum vulgare* (european privet) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Lonicera japonica* - 2006




**Figure 17. Abundance and distribution of *Lonicera japonica* (Japanese honeysuckle) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Lonicera maackii* - 2006




**Figure 18. Abundance and distribution of *Lonicera maackii* (amur honeysuckle) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Lonicera morrowii* - 2006




**Figure 19. Abundance and distribution of *Lonicera morrowii* (morrow's honeysuckle) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Lotus corniculatus* - 2006




**Figure 20. Abundance and distribution of *Lotus corniculatus* (bird's-foot trefoil) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Lysimachia nummularia* - 2006




**Figure 21. Abundance and distribution of *Lysimachia nummularia* (creeping jenny) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Microstegium vimineum* - 2006




**Figure 22. Abundance and distribution of *Microstegium vimineum* (nepalese browntop) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

### *Poa compressa* - 2006




**Figure 23. Abundance and distribution of *Poa compressa* (Canada bluegrass) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Polygonum cuspidatum* - 2006




**Figure 24. Abundance and distribution of *Polygonum cuspidatum* (japanese knotweed) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Robinia pseudoacacia* - 2006




**Figure 25. Abundance and distribution of *Robinia pseudoacacia* (black locust) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Rosa multiflora* - 2006




**Figure 26. Abundance and distribution of *Rosa multiflora* (multiflora rose) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Securigera varia* - 2006




**Figure 27. Abundance and distribution of *Securigera varia* (crownvetch) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Sorghum halepense* - 2006




**Figure 28. Abundance and distribution of *Sorghum halepense* (Johnsongrass) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

### *Spiraea spp* - 2006




**Figure 29. Abundance and distribution of *Spiraea spp* (meadowsweet) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

## *Verbascum thapsus* - 2006




**Figure 30. Abundance and distribution of *Verbascum thapsus* (common mullein) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

### Vinca minor - 2006



**Figure 31. Abundance and distribution of *Vinca minor* (common periwinkle) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

### *Wisteria sinensis* - 2006



**Figure 32. Abundance and distribution of *Wisteria sinensis* (chinese wisteria) at Lincoln Boyhood National Memorial, 2006. Cover classes are as follows: 1=0.1-0.9 m<sup>2</sup>, 2=1-9.9 m<sup>2</sup>, 3=10-49.9 m<sup>2</sup>, 4= 50-99.9 m<sup>2</sup>, 5=100-499.9 m<sup>2</sup>, 6= 499.9-999.9 m<sup>2</sup>, 7=1,000-4,999.9 m<sup>2</sup>, 8=5,000-9,999.9 m<sup>2</sup>, and 9=10,000-14,999.9.**

The NPS has organized its parks with significant natural resources into 32 networks linked by geography and shared natural resource characteristics. HTLN is composed of 15 National Park Service (NPS) units in eight Midwestern states. These parks contain a wide variety of natural and cultural resources including sites focused on commemorating civil war battlefields, Native American heritage, westward expansion, and our U.S. Presidents. The Network is charged with creating inventories of its species and natural features as well as monitoring trends and issues in order to make sound management decisions. Critical inventories help park managers understand the natural resources in their care, while monitoring programs help them understand meaningful change in natural systems and to respond accordingly. The Heartland Network helps to link natural and cultural resources by protecting the habitat of our history.

The I&M program bridges the gap between science and management with a third of its efforts aimed at making information accessible. Each network of parks, such as Heartland, has its own multi-disciplinary team of scientists, support personnel, and seasonal field technicians whose system of online databases and reports make information and research results available to all. Greater efficiency is achieved through shared staff and funding as these core groups of professionals augment work done by individual park staff. Through this type of integration and partnership, network parks are able to accomplish more than a single park could on its own.

The mission of the Heartland Network is to collaboratively develop and conduct scientifically credible inventories and long-term monitoring of park "vital signs" and to distribute this information for use by park staff, partners, and the public, thus enhancing understanding which leads to sound decision making in the preservation of natural resources and cultural history held in trust by the National Park Service.

[www.nature.nps.gov/im/units/htln/](http://www.nature.nps.gov/im/units/htln/)



The U.S. Department of the Interior (DOI) is the nation's principal conservation agency, charged with the mission *"to protect and provide access to our Nation's natural and cultural heritage and honor our trust responsibilities to Indian tribes and our commitments to island communities."* More specifically, Interior protects America's treasures for future generations, provides access to our nation's natural and cultural heritage, offers recreation opportunities, honors its trust responsibilities to American Indians and Alaska Natives and its responsibilities to island communities, conducts scientific research, provides wise stewardship of energy and mineral resources, fosters sound use of land and water resources, and conserves and protects fish and wildlife. The work that we do affects the lives of millions of people; from the family taking a vacation in one of our national parks to the children studying in one of our Indian schools.

NPS D-43, March 2007

**National Park Service**  
**U.S. Department of the Interior**



---

**Natural Resource Program Center**  
1201 Oakridge Drive, Suite 150  
Fort Collins, CO 80525

[www.nps.gov](http://www.nps.gov)

**EXPERIENCE YOUR AMERICA <sup>T</sup>**